



Exam Problem Sheet

The exam consists of 7 problems. You can achieve 55 points in total. The number of points for each problem is marked in brackets (you get 5 points for free). You can find the translation of the problems into Dutch below.

1. [4 Points.] Let R be a ring. The *center* of R is defined as

$$Z(R) = \{a \in R : \forall x \in R : ax = xa\}.$$

Show that $Z(R)$ forms a subring of R .

2. [4+4 Points.] Let R be a ring. For $n \in \mathbb{Z}_{\geq 0}$ and $r \in R$, we define $nr := r + \dots + r$ (n times).

- (a) Suppose R is commutative. Show that

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \quad (*)$$

for all $a, b \in R$ and $n \in \mathbb{Z}_{>0}$. (Hint: you may use that $\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k}$.)

- (b) Show that if conversely Equation (*) holds for all $a, b \in R$ and $n \in \mathbb{Z}_{>0}$ then the ring R is commutative.

3. [4+2 Points.] Let R be a ring, and I, J ideals of R .

- (a) Show that

$$(I + J) \cdot (I \cap J) \subset (I \cdot J) + (J \cdot I). \quad (**)$$

- (b) Show that if $R = \mathbb{Z}$ then equality holds in Equation (**).

4. [3+3+3 Points.]

Let K be a field. The *ring of dual numbers* over K with notation $K[\epsilon]$ consists of expressions of the form $a + b\epsilon$ with $a, b \in K$ which are added and multiplied in the following way:

$$\begin{aligned} (a + b\epsilon) + (c + d\epsilon) &= (a + c) + (b + d)\epsilon, \\ (a + b\epsilon) \cdot (c + d\epsilon) &= (ac) + (ad + bc)\epsilon \end{aligned}$$

(i.e. $\epsilon^2 = 0$), for $a, b, c, d \in K$.

- (a) Prove that $K[\epsilon] \cong K[X]/(X^2)$.
 (b) Prove that $K[\epsilon]$ has exactly *three* ideals.
 (c) Prove that $K[\epsilon]^* \cong K^* \times K^+$ (as groups).

5. [3+3 Points.]

(a) Show that the tangent to the circle

$$S^1 := \{(a, b) \in \mathbb{R}^2 : a^2 + b^2 = 1\}$$

at the point $(a, b) \in S^1$ is given by

$$l_{(a,b)}(X, Y) = 0,$$

where

$$l_{(a,b)}(X, Y) := a(X - a) + b(Y - b) \in \mathbb{R}[X, Y].$$

(b) Let $\mathbb{R}[\epsilon]$ be the ring of dual numbers in Exercise 4. (i.e. $\epsilon^2 = 0$). Define

$$S^1(\mathbb{R}[\epsilon]) := \{(a + s\epsilon, b + t\epsilon) \in \mathbb{R}[\epsilon] \times \mathbb{R}[\epsilon] : (a + s\epsilon)^2 + (b + t\epsilon)^2 = 1\}.$$

Show that:

$$(a + s\epsilon, b + t\epsilon) \in S^1(\mathbb{R}[\epsilon]) \Leftrightarrow l_{(a,b)}(a + s, b + t) = 0.$$

6. [3+3+3 Points.] Let $f : R_1 \rightarrow R_2$ be a (unitary) ring homomorphism between two commutative rings, $I_2 \subset R_2$ an ideal, and $I_1 := f^{-1}(I_2) \subset R_1$.

(a) Prove: I_1 is an ideal in R_1 , and R_1/I_1 is isomorphic to a subring of R_2/I_2 .

(b) Prove: if I_2 is prime in R_2 then I_1 is prime in R_1 .

(c) Give an example which shows that (b) can be wrong if both occurrences of ‘prime’ are replaced by ‘maximal’.

7. [2+2+2+2 Points.]

(a) Show that $X^n + 2$ is irreducible in $\mathbb{Z}[X]$ for all $n \in \mathbb{Z}_{\geq 0}$.

(b) Show that $Y^n - X$ is irreducible in $K[X, Y]$ (K a field) for all $n \in \mathbb{Z}_{\geq 0}$.

(c) Give the example of an irreducible polynomial $f \in \mathbb{Z}[X]$ with the property that $f(X^2)$ is *not* irreducible.

(d) Let $f \in \mathbb{Z}[X]$ be a monic Eisenstein polynomial. Show that $f(X^2)$ is irreducible in $\mathbb{Z}[X]$.

Dutch Translation

1. [4 Punten.] Zij R een ring. Het *centrum* van R is

$$Z(R) = \{a \in R : \forall x \in R : ax = xa\}.$$

Bewijs dat dit een deelring van R is.

2. [4+4 Punten.] Laat R een ring zijn. Voor $n \in \mathbb{Z}_{\geq 0}$ en $r \in R$ definiëren we $nr := r + \dots + r$ (n keer).

(a) Stel R is commutatief. Bewijs dat

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \quad (\star)$$

voor alle $a, b \in R$ en $n \in \mathbb{Z}_{>0}$. (Aanwijzing: Je mag gebruiken dat $\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k}$.)

(b) Bewijs omgekeerd, dat als (\star) voor alle $a, b \in R$ en $n \in \mathbb{Z}_{>0}$ geldt, de ring R commutatief is.

3. [4+2 Punten.]

Zij R een ring, en I, J idealen van R .

(a) Bewijs

$$(I + J) \cdot (I \cap J) \subset (I \cdot J) + (J \cdot I). \quad (\star\star)$$

8

(b) Bewijs dat gelijkheid in $(\star\star)$ geldt als $R = \mathbb{Z}$.

4. [3+3+3 Punten.]

Zij K een lichaam. De ring van de duale getallen over K , notatie: $K[\epsilon]$, bestaat uit de uitdrukkingen $a + b\epsilon$, met $a, b \in K$, die als volgt opgeteld en vermenigvuldigd worden:

$$\begin{aligned} (a + b\epsilon) + (c + d\epsilon) &= (a + c) + (b + d)\epsilon, \\ (a + b\epsilon) \cdot (c + d\epsilon) &= (ac) + (ad + bc)\epsilon \end{aligned}$$

(dus $\epsilon^2 = 0$), voor $a, b, c, d \in K$.

(a) Bewijs: $K[\epsilon] \cong K[X]/(X^2)$.

— (b) Bewijs dat $K[\epsilon]$ precies drie idealen heeft.

— (c) Bewijs: $K[\epsilon]^* \cong K^* \times K^+$ (als groepen).

→ 5. [3+3 Punten.]

(a) Ga na dat de raaklijn aan de cirkel

het

$$S^1 := \{(a, b) \in \mathbb{R}^2 : a^2 + b^2 = 1\}$$

in de punt $(a, b) \in S^1$ gegeven is door

$$l_{(a,b)}(X, Y) = 0,$$

waarbij

$$l_{(a,b)}(X, Y) := a(X - a) + b(Y - b) \in \mathbb{R}[X, Y].$$

(b) Zij $\mathbb{R}[\epsilon]$ de ring van de duale getallen in Exercise 4. (d.w.z. $\epsilon^2 = 0$). Definieer

$$S^1(\mathbb{R}[\epsilon]) := \{(a + s\epsilon, b + t\epsilon) \in \mathbb{R}[\epsilon] \times \mathbb{R}[\epsilon] : (a + s\epsilon)^2 + (b + t\epsilon)^2 = 1\}.$$

Laat zien dat :

$$(a + s\epsilon, b + t\epsilon) \in S^1(\mathbb{R}[\epsilon]) \Leftrightarrow l_{(a,b)}(a + s, b + t) = 0.$$

6. [3+3+3 Punten.] Zij $f : R_1 \rightarrow R_2$ een (unitair) homomorfisme van commutatieve ringen, $I_2 \subset R_2$ een ideaal, en $I_1 := f^{-1}(I_2) \subset R_1$.

— (a) Bewijs: I_1 is een ideaal in R_1 , en R_1/I_1 is isomorf met een deelring van R_2/I_2 .

— (b) Bewijs: als I_2 priem is in R_2 dan is I_1 priem in R_1 .

— (c) Laat an de hand van een voorbeeld zien dat (b) fout can zijn als 'priem' beide malen vervangen wordt door 'maximaal'.

7. [2+2+2+2 Punten.]

(a) Bewijs dat $X^n + 2$ irreducibel in $\mathbb{Z}[X]$ is voor alle $n \in \mathbb{Z}_{>0}$.

— (b) Bewijs dat $Y^n - X$ irreducibel is in $K[X, Y]$ (K een lichaam) voor alle $n \in \mathbb{Z}_{>0}$.

— (c) Vind een voorbeeld van een irreducibel polynoom $f \in \mathbb{Z}[X]$ met de eigenschap dat $f(X^2)$ niet irreducibel is.

(d) Laat $f \in \mathbb{Z}[X]$ een monisch Eisensteinpolynoom zijn. Bewijs dat $f(X^2)$ irreducibel in $\mathbb{Z}[X]$ is.