

Analysis on Manifolds (WBMA013-05)– Final Exam

Tuesday 26 January 2021, 8:30h–12:30h

This exam consists of 3 problems.

Usage of the theory and examples from the lecture notes is allowed with the only exception of the results of Exercise 4.1.13 from the lecture notes. Give a precise reference to the theory and/or exercises you use for solving the problems.

You get 10 points for free.

Problem 1. (9 + 15 + 6 = 30 points)

Let $f(x, y, z) = x^2 + y^2 + z^2$ and let $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be the map

$$\sigma(u, v) = \frac{1}{u^2 + v^2 + 1} (2u, 2v, u^2 + v^2 - 1). \quad (1)$$

- (a) If $X = z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z}$, compute $\iota_X df$.
- (b) Verify that $d(f \circ \sigma) = \sigma^* df$ by computing separately both terms.
- (c) Does $du \wedge \sigma^* df$ define a volume form on \mathbb{R}^2 ? If so, is it positively oriented with respect to the standard euclidean basis? Justify your answer.

Problem 2. (6 + 6 + 6 + 6 + 6 = 30 points)

Recall that we can identify the space $\text{Mat}(2, \mathbb{R})$ of 2×2 -matrices with \mathbb{R}^4 by associating the matrix $X = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}$ with the point $(x_{11}, x_{12}, x_{21}, x_{22}) \in \mathbb{R}^4$.

- (a) Show that the set

$$\text{SL}(2, \mathbb{R}) = \{A \in \text{Mat}(2, \mathbb{R}) \mid \det A = 1\}$$

is a 3-dimensional smooth submanifold of $\text{Mat}(2, \mathbb{R})$.

Hint: use the identification between matrices and \mathbb{R}^4 .

- (b) Let $e \in \text{Mat}(2, \mathbb{R})$ denote the identity matrix. Show that

$$T_e \text{SL}(2, \mathbb{R}) = \{A \in \text{Mat}(2, \mathbb{R}) \mid \text{tr} A = 0\},$$

where $\text{tr} A$ denotes the matrix trace, i.e., the sum of the diagonal entries of A .

- (c) Let $\iota : \text{SL}(2, \mathbb{R}) \rightarrow \text{SL}(2, \mathbb{R})$ be the map $\iota(A) = A^{-1}$. Show that ι is smooth.
- (d) Show that $d\iota_e : T_e \text{SL}(2, \mathbb{R}) \rightarrow T_e \text{SL}(2, \mathbb{R})$ is given by $d\iota_e(A) = -A$.
- (e) Show that $\text{SL}(2, \mathbb{R})$ is a Lie group and give its Lie algebra.

Problem 3. (6 + 6 + 10 + 8 = 30 points)

In this problem we are going to prove the following fixed point theorem.

Theorem 1. Let $D_n := \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$ denote the closed unit disk in \mathbb{R}^n . Any smooth map $g : D_n \rightarrow D_n$ has a fixed point, that is, $\exists p \in D_n$ such that $g(p) = p$.

We will proceed by first showing another result.

Theorem 2. Let N be a compact n -dimensional submanifold of \mathbb{R}^n with non-empty boundary ∂N . Then, there is no differentiable map $f : N \rightarrow \partial N$ for which every boundary point is a fixed point, that is, for which $f(p) = p$ for all $p \in \partial N$.

Let $\Omega = dx^1 \wedge \cdots \wedge dx^n$ denote the standard volume form on N , that is, the restriction of the standard volume form on \mathbb{R}^n to N , and X be an outward-pointing vector field on ∂N .

- (a) Show that $\omega = \iota_X \Omega$ is a closed non-vanishing form on ∂N .
- (b) Show that if there is f such that $f(p) = p$ for all $p \in \partial N$, then $f^* \omega$ is closed.
- (c) Prove Theorem 2.
Hint: use integration to get a contradiction.
- (d) Prove Theorem 1.
Hint: by contradiction, consider $f(p) = \frac{p-g(p)}{\|p-g(p)\|} \cdots$