

Probability Theory
2016/17 Semester IIb
Instructor: Daniel Valesin
Final Exam
22/6/2017
Duration: 3 hours

Name: _____
Student number: _____

This exam contains 9 pages (including this cover page) and 7 problems. Enter all requested information on the top of this page.

Your answers should be written in this booklet. Avoid handing in extra paper.

You are allowed to have two hand-written sheets of paper and a calculator.

You are required to show your work on each problem.

Do not write on the table below.

Problem	Points	Score
1	14	
2	14	
3	14	
4	14	
5	10	
6	14	
7	10	
Total:	90	

1. (a) (7 points) In how many ways can we distribute m indistinguishable candies to k distinguishable children, assuming $m \geq k$ and each child gets at least one candy?
- (b) (7 points) Half the lamps produced by a factory are of good quality and half of bad quality. The lifetime (in months) of a good-quality lamp is an exponential random variable with parameter $\beta = 2$, and the lifetime of a bad-quality lamp is an exponential random variable with $\beta = 1$. Assume that you pick a lamp at random in the factory, install it and, after t months, it is still working. Let $p(t)$ be the probability that you now attribute to having picked a good-quality lamp. Find the limit of $p(t)$ as $t \rightarrow \infty$.

2. (a) (7 points) The discrete random variables X and Y are jointly distributed as follows. X follows the $\text{Poisson}(\lambda)$ distribution, and

$$f_{Y|X}(y|x) = \binom{x}{y} p^y (1-p)^{x-y}, \quad x \in \{0, 1, \dots\}, y \in \{0, \dots, x\},$$

where $p \in (0, 1)$. Show that the distribution of Y is $\text{Poisson}(\lambda p)$.

- (b) (7 points) The continuous random variables Z and W are independent, with Z following the exponential distribution with parameter 1 and W following the (continuous) uniform distribution on $(0, 1)$. Find $\mathbb{P}(Z < W < 3Z)$.

3. Let A be a set with n elements. Assume that we choose a subset of A at random according to the rule that the probability that a certain subset A' is chosen is proportional to the number of elements of A' . Let X be the number of elements of the subset we choose.

(a) (7 points) Find the probability mass function of X .

(b) (7 points) Prove that

$$M_X(t) = e^t \left(\frac{e^t + 1}{2} \right)^{n-1}, \quad t \in \mathbb{R}.$$

4. (a) (7 points) A device has lifetime denoted by T , which follows an exponential distribution with parameter $\beta = 1.5$. The device has value $V = 5$ if it fails before $t = 3$; otherwise, it has value $V = 2T$. Find the cumulative distribution function of V .
- (b) (7 points) Let X_1, X_2, \dots, X_n be independent random variables, all following an exponential distribution with parameter β . Let

$$Y = \max\{X_1, \dots, X_n\},$$

that is, Y is equal to the largest among the values X_1, \dots, X_n . Find the cumulative distribution function of Y .

5. Let X be a random variable with expectation μ_X and variance σ_X^2 , and Y a random variable with expectation μ_Y and variance σ_Y^2 . Let $\rho_{X,Y}$ be the correlation between X and Y . Express the following quantities in terms of μ_X , μ_Y , σ_X , σ_Y , and $\rho_{X,Y}$.
- (a) (5 points) $\text{Var}(X - 3Y)$;
 - (b) (5 points) $\mathbb{E}((3X - 5)(2Y + 1))$.

6. (a) (7 points) Let X_1, X_2, \dots be independent and identically distributed random variables with finite expectation and variance. Let $Z_i = X_i + X_{i+1}$, for $i = 1, 2, \dots$. Show that $\bar{Z}_n = \frac{1}{n} \sum_{i=1}^n Z_i$ converges in probability to a constant, and identify this constant.
- (b) (7 points) Let Y_1, Y_2, \dots be independent and identically distributed *positive* random variables with finite expectation μ and finite variance σ^2 . For each natural number n , let W_n be the largest value of k for which the following inequality holds:

$$\sum_{i=1}^k Y_i \leq n.$$

Show that W_n/n converges in probability to $1/\mu$.

Hint. Use the facts that $\{W_n > x\} \subseteq \{\sum_{i=1}^{\lfloor x \rfloor} Y_i < n\}$, $\{W_n < x\} \subseteq \{\sum_{i=1}^{\lceil x \rceil} Y_i > n\}$.

7. (10 points) At each time instant $t = 0, 1, 2, \dots$, a particle occupies a position in the set $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$. Assume that the initial position is 0 and, from time t to time $t + 1$, the particle moves:
- one unit to the left with probability $1/3$;
 - one unit to the right with probability $1/6$;
 - two units to the right with probability $1/2$.

Find a value k such that the probability that the particle is to the left of k at time 10000 is approximately 70%.

If you don't have a calculator, you may use the approximation: $\sqrt{65} \approx 8.06$.

Note that a table for the cumulative distribution function of the standard normal distribution is provided on page 2.