

Linear Algebra I

09/10/2023

You are **NOT** allowed to use any type of calculators.

1 Systems of linear equations

(1 + 6 + 3 + (1 + 6 + 3) = 20 pts)

Let a be a scalar. Consider the following system of linear equations in the unknowns x , y , and z :

$$\begin{aligned}x + 2y - 3z &= 4 \\3x - y + 5z &= -2 \\4x + y + (a^2 - 14)z &= a + 6.\end{aligned}$$

- Write down the corresponding augmented matrix .
- By performing elementary row operations, put the augmented matrix into row echelon form.
- Determine all values of a so that the system is consistent.
- For $a = 5$,
 - determine the *lead* and *free* variables.
 - put the augmented matrix into *reduced* row echelon form by performing elementary row operations.
 - find the solution set.

2 Matrix multiplication

(5 + 10 = 15 pts)

Prove or disprove the statements: For all $A \in \mathbb{F}^{p \times q}$ and $B \in \mathbb{F}^{q \times p}$,

- $\det(AB) = \det(BA)$.
- $\operatorname{tr}(AB) = \operatorname{tr}(BA)$. (For $M \in \mathbb{F}^{m \times m}$, $\operatorname{tr}(M) := \sum_{k=1}^m [M]_{kk}$.)

3 Determinants

(10 + 15 = 25 pts)

Let $M(n) \in \mathbb{R}^{n \times n}$ be given by

$$[M(n)]_{ij} = \begin{cases} 1 & \text{if } |i - j| \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

For instance,

$$M(5) = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

- (a) Compute the determinant of $M(n)$ for $n \in \{1, 2, 3\}$.
- (b) Find real numbers a, b such that $\det(M(n+2)) = a \det(M(n+1)) + b \det(M(n))$ for all $n \geq 1$.

4 Nonsingular matrices

(15 + 15 = 30 pts)

Let a, b, c, d be scalars and $M \in \mathbb{F}^{m \times m}$. Consider the matrix

$$N = \begin{bmatrix} aM & bM \\ cM & dM \end{bmatrix}.$$

- (a) Show that N is nonsingular if and only if $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and M are nonsingular.
- (b) Suppose that $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and M are nonsingular. Find the inverse of N .

10 pts free